

# Changes Late Blight Fungicide Table

Huub Schepers & Bert Evenhuis



# EuroBlight Table

- Changes since Workshop Brasov 2015
  - Accuracy of some first registration dates should be checked: no amendments
  - Link to Early Blight Table should be clearer: done
  - No host resistance elicitors were tested
  - Where there is suspicion of a discrepancy between a fungicide's rating and its current efficacy, advisors need to report this: no cases received
  - Links to Latin-Blight, USA-Blight and Asia-Blight on website: yes in Newsletter but no links yet
  - Protocols should be shared: partly done

---

# EuroBlight Late Blight Fungicide Table

Based on EuroBlight leaf trials:

\* (zoxamide + dimethomorph) + fluazinam: 4.6

Based on EuroBlight tuber trials:

\* no changes

Based on company dossier & experts:

\* mandipropamid + cymoxanil: qualitative ratings added

# Summary discussion 17 May 2017

## Control Strategies

### ■ Best Practices

- Update 11 EuroBlight BPs: Faye, Ruairidh, Huub
- Adapt TizonLatino BPs; Jorge, Ivette

### ■ Protocols

- Upload all available protocols on website
- Update protocols: ?

### ■ Statement agrochemicals

- Express need of network for broad spectrum of fungicides (and herbicides) as tools for IPM
- Resistance management: 4 MoA per disease/crop
- Also for specific use (new growth, tuber etc.)